

Transfinite diameter of Bernstein sets

Dimitri Jordan Kenne

Faculty of Mathematics and Computer Sciences, Jagiellonian University, Krakow, Poland

keywords: Transfinite diameter, Bernstein set, Markov set

Abstract:

Let $K \subset \mathbb{C}^n$ be a compact set satisfying the following Bernstein inequality: for any $m \in \{0, \dots, n\}$ and for any n -variate polynomial P of degree $\deg(P)$ we have

$$\max_{z \in K} \left| \frac{\partial P}{\partial z_m}(z) \right| \leq M \deg(P) \max_{z \in K} |P(z)|. \quad (1)$$

for some constant $M := M(K) > 0$ depending only on K . We show that the transfinite diameter of K , denoted $\delta(K)$, verifies the following lower estimate

$$\delta(K) \geq \frac{1}{nM}, \quad (2)$$

which is optimal in the one-dimensional case.

References

- L. Białas-Cieź and M. Jedrzejowski. Transfinite diameter of Bernstein sets in \mathbb{C}^n . *J Inequal Appl*, 2002 (3):680202, 2002.
- D. J. Kenne. Transfinite diameter of bernstein sets. *Preprint*, 2022.
- F. Toókos and V. Totik. Markov inequality and Green functions. *Rend. Circ. Mat. Palermo math*, 2: 91–102, 2005.
- Ö. YAZICI. A note on the transfinite diameter of Bernstein sets. *Turkish Journal of Mathematics*, 46(7): 2761–2765, 2022.